

Relativistic effects in parity-violating electron-deuteron scattering and strangeness in the nucleon.

Grigorios I. Poulis^{a,b}

(a) *Department of Physics and Mathematical Physics
and Institute for Theoretical Physics*

*University of Adelaide, 5005 Adelaide SA, Australia*¹
and

(b) *NIKHEF, P.O. Box 41882, 1009 DB Amsterdam, The Netherlands*

ABSTRACT

Deuteron electrodisintegration is described in a relativistic PWIA framework. The reduction to the factorized PWIA, and the effect thereof on the observables accessible in parity-violating scattering, is discussed. The extent to which such relativistic effects may affect the extraction of strangeness nucleon form factors from measurements of the PV asymmetry, is investigated.

Parity-violating (PV) inclusive scattering of polarized electrons off unpolarized protons or nuclei provides a tool for extracting the hitherto unknown strangeness nucleon form factors $G_{E,M}^s$. The SAMPLE experiment at MIT-Bates [1], a $Q^2=0.1$ (GeV/c)², backward angles measurement of the $p(\vec{e}, e')p$ asymmetry, is focusing on G_M^s . However, strong correlation with the neutral current, axial, isovector form factor, $\tilde{G}_A^{T=1}$, whose radiative corrections have a large theoretical uncertainty in the Standard Model, places constraints on the effectiveness of the SAMPLE measurement. It has been suggested [2] that a complimentary measurement of the PV asymmetry in quasielastic electron-deuteron scattering, where this correlation is suppressed by a factor of 5, may, in conjunction with the proton measurement, allow one to determine both G_M^s and $\tilde{G}_A^{T=1}$. A second phase of the SAMPLE experiment, as well as a TJNAF (ex-CEBAF) experiment will perform such measurements on the deuteron [1]. Hadjimichael et al. [2] have estimated the unavoidable theoretical uncertainty that nuclear dynamics introduces in the modeling of the ${}^2H(\vec{e}, e')np$ reaction and found that relativistic, rather than rescattering effects, incorporated via factorized (F) PWIA, are the main source of model dependence in the PV asymmetry for quasifree kinematics at $q > 0.7$ GeV/c, especially at forward angles (6% effects). In this work we treat the ${}^2H(\vec{e}, e')np$ reaction in relativistic (R) PWIA, where not only the single-nucleon tensor (as in FPWIA), but also the d-n-p vertex is treated relativistically. To the extent that for quasifree kinematics the two scales governing the relativistic effects in this reaction, q/m and $|\vec{k}|/m$, decouple (Fig. 1), one expects that, since $|\vec{k}|$ is restricted by the small (≈ 60 MeV/c) deuteron Fermi momentum, at high q values the FPWIA employed in [2] (although it does not treat relativistic effects associated with the $|\vec{k}|/m$ scale properly) should not differ significantly from the RPWIA.

¹present address

The purpose of this work is to test this conjecture. The kinematics (deuteron rest frame) are described in Fig. 1; $q^\mu = (\omega, \vec{q})$, $d^\mu = (M_d, \vec{0})$, $k^\mu = (E_k, \vec{k})$, $n^\mu = (E_n, -\vec{k})$, $p^\mu = q^\mu + k^\mu = (E_p, \vec{p})$, with $d^2 = M_d^2$, $p^2 = n^2 = m^2$, $k^2 < m^2$ (off-shell struck nucleon). In the (semirelativistic) FPWIA the hadronic tensor $W_F^{\mu\nu}$ factorizes as a non-relativistic (NR) momentum distribution $\rho(|\vec{k}|)$ times a relativistic off-shell single-nucleon tensor $\mathcal{W}_F^{\mu\nu}$:




Fig. 1. PWIA hadronic tensor.

$$W_F^{\mu\nu} = (2\pi)^3 M_d \rho(|\vec{k}|) \mathcal{W}_F^{\mu\nu}, \quad \text{with } \mathcal{W}_F^{\mu\nu} = \text{Tr} \left[(\vec{k} + m) \bar{\Gamma}^\mu (\not{p} + m) \Gamma^\nu \right] (2m)^{-2}. \quad (1)$$

Here Γ_μ is the electromagnetic (E/M) or neutral current (NC) nucleon vertex, $\bar{\Gamma}_\mu = \gamma_0 \Gamma_\mu^\dagger \gamma_0$, and the de Forest [3] prescription is used – i.e., the $(\vec{k} + m)$ projector results from treating the struck nucleon as “would-be-on-shell”, using $\vec{k}^\mu = (\vec{E}_k, \vec{k})$, where $\vec{E}_k^2 = E_n^2 = m^2 + \vec{k}^2$. In RPWIA, on the other hand, one uses the covariant d-n-p vertex $\mathcal{X}^\mu = [A\gamma^\mu + B\delta^\mu + (m - \not{k})(F\gamma^\mu + G\delta^\mu)]$, where $\delta = (n - k)/2$, and the vertex functions A, B, F, G can be written in terms of the $^3S_1$, (u) $^3D_1$, (w) , $^3P_1$, (v_t) , $^1P_1$ (v_s) relativistic deuteron wavefunctions. In this work we use the wavefunctions of Ref. [4], obtained from the spectator equation with pseudoscalar πNN coupling, which enhances the role of the P states. The hadronic tensor reads in PWIA

$$W_R^{\mu\nu} = \frac{1}{3(k^2 - m^2)^2} \text{Tr} \left[\frac{\not{p} + \beta}{2m} (\not{k} + m) \bar{\Gamma}^\mu \frac{\not{p} + m}{2m} \Gamma^\nu (\not{k} + m) \right], \quad (2)$$

where $(\not{p} + \beta) = (-g_{\rho\lambda} + d_\rho d_\lambda / M_d^2) \mathcal{X}^\lambda (\not{p} - m) \bar{\mathcal{X}}^\rho$. In order to reduce the above, *non*-factorized RPWIA to the FPWIA of Eq. (1), we split the struck nucleon propagator into + and – energy contributions: $\not{k} + m = f_+(\not{k}_+ + m) + f_-(\not{k}_- - m)$, where (in the target rest frame) $f_\pm = (E_k \pm \vec{E}_k) / 2\vec{E}_k$, $k_+ = \vec{k}$, $k_- = n$. We then find that a *weak*, density-matrix type of factorization still holds (details will appear elsewhere)

$$W_R^{\mu\nu} = (2\pi)^3 M_d \sum_{\text{spins}} [\rho_{++} \mathcal{W}_{++}^{\mu\nu} + \rho_{+-} \mathcal{W}_{+-}^{\mu\nu} + \rho_{-+} \mathcal{W}_{-+}^{\mu\nu} + \rho_{--} \mathcal{W}_{--}^{\mu\nu}], \quad (3)$$

where, in the shorthand notation $\chi_+^s = u(\vec{k}, s)$ and $\chi_-^s = v(-\vec{k}, s)$, the generalized, relativistic “momentum distributions” and “single-nucleon” tensors are defined as $\rho_{ab}^{s,r} \sim f_a f_b (k^2 - m^2)^{-2} \text{Tr}[\chi_a^s \bar{\chi}_b^r (\not{p} + \beta)]$ and $W_{ab}^{\mu\nu} = \text{Tr}[\chi_b^r \bar{\chi}_a^s \bar{\Gamma}^\mu (\not{p} + m) \Gamma^\nu] / 2m$, respectively. Evaluating the traces we find that the ++ and -- spectral functions are diagonal in spin space, $\rho_{\pm\pm}^{s,r} \sim \delta_{s,r} f_\pm^2 (k^2 - m^2)^{-2} [(k_\pm \cdot \mathcal{P}) / m \pm \beta]$, as well, which then allows one to identify $\delta_{r,s} \mathcal{W}_{++}^{\mu\nu} = \mathcal{W}_F^{\mu\nu}$, [cf. Eq. (1)]. Furthermore, by taking the NR limit of ρ_{++} (which is oblivious to the “hard” scale q/m), i.e., by ignoring the P states and expanding in $|\vec{k}|/m$, we find $\rho_{++} \rightarrow [u^2(|\vec{k}|) + w^2(|\vec{k}|)] / 4\pi = \rho(|\vec{k}|)$, the nonrelativistic momentum distribution in Eq. (1). Thus, the widely used de Forest FPWIA may be obtained from RPWIA in a two-step reduction: (a) truncation to the ++ components of the struck nucleon, (b) additionally, taking the nonrelativistic limit of the ρ_{++} spectral function.




Fig. 2. E/M responses at $q = 1$ GeV/c.

relativistic effects in inclusive, quasielastic scattering are incorporated in the FPWIA, which should indeed be used at large q values, where a comparison with NR treatments shows sizable effects, as mentioned above. Relativistic effects associated with negative energy components of the off-shell struck nucleon and with the target wavefunction are (in the target rest frame!) negligible for such kinematics. Since we are primarily interested in the interplay between the relativistic effects *not* incorporated in the FPWIA and the extraction of strangeness, we show in Fig. 3 the PV asymmetry for a SAMPLE-type experiment, with three values of strangeness magnetic moment μ_s [1]. Interestingly enough, it turns out that relativistic effects beyond FPWIA in the quasielastic asymmetry are maximal for $200 < q < 300$ MeV/c and backward angles, e.g, at the SAMPLE kinematics, where they induce a 0.45% effect, while $A = A_0(1 - 0.055\mu_s)$. The extraction of strangeness on the quasielastic peak is nevertheless not obscured by such residual effects unless μ_s is quite small ($|\mu_s| < 0.1$).




Fig. 3. Asymmetry at $q = 300$ MeV/c. Lower curves: RPWIA. Upper curves: FPWIA.

[1] E.J. Beise et al., archive: nucl-ex/9602001; also E.J. Beise, these proceedings.

[2] E. Hadjimichael, G.I. Poulis and T.W. Donnelly, Phys. Rev. C45, 2666 (1992).

[3] T. de Forest, Nucl. Phys. A392, 232 (1983).

[4] W.W. Buck and F. Gross, Phys. Rev. C20, 2361, (1979).